


## Soapstock in Ruminant Diets

by Davy R. Brown, Ph.D.  
Ruminant Nutritionist

Net energy (NE) intake by feedlot cattle is one of the most important, if not the most important, dietary factors influencing weight gain. Increasing the energy content of the diet by adding fat (1-4% DM basis) results in increased NE intake and rate of gain provided DM intake, fat digestibility, and fiber digestibility are not substantially depressed. Tallow supplementation often results in reduced DMI but with minimal effects on fiber digestibility, resulting in a net increase in energy intake and dietary NE content. The degree to which fat may influence DM intake and nutrient digestibility varies with supplementation level, degree of saturation, and whether or not the fat is protected from the rumen. Unsaturated fats are highly subject to saturation by rumen bacteria through the process of biohydrogenation. This is the primary reason why beef fat and muscle have a higher saturated fat content compared with pork and poultry. Protecting dietary unsaturated fats from biohydrogenation has resulted in increased DM intake, NDF, ADF, and fat digestibility, and higher unsaturated fat levels in blood and tissue when compared with cattle fed diets containing tallow or unprotected vegetable oils (Trenkle et al., 1995; Rule et al., 1994; Pantoja et al., 1994; Doreau et al., 1993; Brandt and Anderson, 1990).

Ruminal availability, and thus biohydrogenation of unsaturated fat may be influenced by grain processing (if oil is in the seed) or by saponification of supplemental fat (i.e. soapstock). Soapstocks (SS) are saponified fatty acids (FA) formed when free fatty acids and divalent cations (usually



calcium) are combined to form a protected fat that is relatively inert in the rumen (Grummer, 1988) but dissociates in the acidic environment of the duodenum (first part of small intestine) during digestion. Dissociation of fat and Ca is important for proper digestion and absorption of soapstocks. Soapstocks can be formed from most fats and oils including (in general order of increasing maturation) tallow, palm oil, cottonseed oil, corn oil, soybean oil, rapeseed oil, and canola oil. Dissociation and digestibility of SS in the rumen and total tract can vary with degree of FA saturation and pH (Sukhija and Palmquist, 1990).

Recent research showed unsaturated soaps were generally more digestible (Enjalbert et al., 1997) and tended to dissociate at a higher pH compared with those that were more saturated. The unique properties of unsaturated SS allowed for DM intake that was similar to tallow supplemented diets but did not negatively influence digestion and metabolism as would be expected with diets containing unprotected fats (Pantoja et al., 1994; Wu et al., 1991). Studies generally agreed that digestibility of major nutrients such as organic matter, starch, nitrogen, and energy were similar while NDF and ADF digestion was improved when unsaturated SS was supplemented (2-6% of diet DM) compared with tallow and vegetable fat supplemented at the same level (Doreau et al., 1993; Zinn, 1992; Ngidi et al., 1990). These findings were consistent with those by White (1958) who observed low fiber digestion was completely reversed when dietary Ca was increased from .3 to .9% in diets supplemented with 5% corn oil. This also suggests when supplying supplemental fat as SS additional dietary Ca may not be needed for improved fiber digestion to occur

(Zinn and Shen, 1996). In agreement with this are the trials reviewed here where improved fiber digestion occurred with unsaturated SS supplementation but with no additional Ca provided.

Even though feeding SS has been associated with improved lilt and fiber digestion, improvements in feedlot performance by steers fed SS compared with those fed tallow or yellow-grease have not been demonstrated (Fluharty and Loerch, 1997; Bock et al., 1991; Ngidi et al., 1990). Nonetheless, higher NE content and improved feedlot performance occurred when diets containing 2-4% SS was fed compared with control diets containing no added fat. Compared with unprotected fat, SS appears to support similar levels of feedlot performance but without the tendency to depress DM intake and fiber digestion. When SS from tallow (i.e. saturated SS) was fed (Jenkins and Palmquist, 1984) ruminal fiber digestibility reached normal levels (e.g. control level) but total fat digestion was reduced. This would be expected since SS is inert in the rumen

and digestibility of saturated fat is lower than that for unsaturated. Unsaturated SS appears to be of benefit since it is relatively inert in the rumen compared with unprotected oils, and is more digestible in the small intestine than is saturated fat and saturated SS. Lastly, if SS is fed as the sole fat source it should probably be supplemented at a level of less than 6% of the total diet DM.

## Literature Cited

- Brandt, R.T. and S.J. Anderson. 1990. Supplemental fat affects feedlot performance and carcass traits of finishing yearling steers and estimated diet NE value. *J. Anim. Sci.* 68:2208-2216.
- Bock, B.J. D.L. Harmon. R.T. Brandt, J.E. Schneider. 1991. Fat source and calcium level effects on finishing steer performance, digestion, and metabolism. *J. Anim. Sci.* 69:2211-2224.
- Enjalbert, F., M.C. Nicol, C. Bayourthe, M. Vernay and R. Moncoulon. 1997. Effects of dietary calcium soaps on digestion, milk composition, and physical properties of butter. *J. Dairy Res.* 64:181-195.
- Doreau, M., A. Ferlay, and Y. Elmeddah. 1993. Organic matter and nitrogen digestion by dairy cows fed calcium salts of rapeseed oil fatty acids or rapeseed oil. *J. Anim. Sci.* 71:499-504.
- Fluharty, F.L. and S.C. Loerch 1997. Effects of source and supplemental fat and protein on performance of newly arrived feedlot steers. *J. Anim. Sci.* 75:2308-2316.
- Grummer, R.R. 1988. Influenced of prilled fat and calcium salt of palm oil fatty acids on ruminal fermentation and nutrient digestibility. *J. Dairy Sci.* 71:117-123.
- Jenkins, T.C. and D.L. Palmquist. 1984. Effects of fatty acids or calcium soaps on rumen and total nutrient digestibility of dairy rations. *J. Dairy. Sci.* 67:978-986.
- Ngidi, M.E, S.C. Welch, F.L. Fluharty, and D.L. Palmquist. 1990. Effects of calcium soaps of long-chain fatty acids on feedlot performance, carcass, characteristics, and ruminal metabolism of steers. *J. Anim. Sci.* 68:2555-2565.
- Ohajuruka, O.A., Z.G. Wu, and D.L. Palmquist. 1991. Ruminal metabolism, fiber, and protein digestion by lactating cows fed calcium soaps or animal-vegetable fat. *J. Anim. Sci.* 74:2601-2609
- Pantoja, J., J.L. Firkins, M.L. Eastridge, and B.L. Hull. 1994. Effects of fat saturation and source of fiber on site of nutrient digestion and milk production by lactating cows. *J. Dairy Sci.* 77:2341-2356.
- Sukhija, P.S. and D.L. Palmquist. 1990. Dissociation of calcium soaps of long-chain fatty acids in rumen fluid. *J. Dairy. Sci.* 73:1784-1787.
- White, T.W., R.B. Grainger, F.H. Baker, and J.W. Stroud. 1958. Effect of supplemental fat on digestion and rum final calcium requirement of sheep. *J. Anim. Sci.* 17:797.
- Wu, Z., O.A. Ohajuruka, D.L. Palmquist. 1991. Ruminal synthesis, biohydrogenation, and digestibility of fatty acids by dairy cows. *J. Dairy Sci.* 74:3025-3034.
- Zinn, R.A. and Y. Shen. 1996. Interaction of dietary calcium and supplemental fat oil digestive function and growth performance in feedlot steers. *J. Anim. Sci.* 74:2303-2309.
- Zinn, R.A. 1992. Comparative feeding value of supplemental fat in steam-flaked corn- and steam-flaked wheat-based finishing diets for feedlot steers. *J. Anim. Sci.* 70:2959-2969.

 **MIX 30**<sup>™</sup>  
THE HIGH ENERGY LIQUID FEED

